

Original Article

Gingivitis and periodontitis as a risk factor for stroke: A case-control study in the Iranian population

Maryam Alsadat Hashemipour¹, Afsaneh Jahanshahi Afshar², Roya Borna², Behnas Seddighi³, Amin Motamedi⁴

¹Member of Kerman Neuroscience Research Center, Member of Kerman Oral and Dental Diseases Research Center, and Department of Oral Medicine, ²Member of Kerman Social Determinants on Oral Health Research Center, ³Department of Neurology, ⁴Member of Kerman Neuroscience Research Center, and Member of Kerman Oral and Dental Diseases Research Center, Department of Periodontology, Faculty of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

ABSTRACT

Background: Periodontitis and gingivitis are one of the most infectious diseases in human. Several studies have been carried out on dependence of periodontitis and stroke. The aim of this study was to investigate the gingivitis and periodontitis as a risk factor for stroke in Iranian population.

Materials and Methods: A case-control study was conducted on 100 patients suffering from stroke as case group, and 100 hospitalized patients as control group. The case group included 42 males and 58 females, and in control group there were 44 males and 56 females. Using a University of North Carolina-15 manual probe, the clinical attachment level, the distance between the cemento-enamel junction and the probed base of the periodontal pocket, were recorded by gingival and periodontal indexes. The data were analyzed by multiple logistic regressions, Chi-square test, Fisher's test, t-test, Man Whitney, and SPSS 11.5 software program. $P < 0.05$ was considered as significant.

Results: The case group included 42 males and 58 females, and in control group 44 males and 56 females were included. In this investigation, the average of gingival index in men and women of case group was 1.22 ± 0.55 and 1.31 ± 0.55 , respectively. This study showed that the average of gingival index in case group was more than control group. Periodontal index in both groups in men was more than women. The moderate and severe periodontitis in case group were more than that of control group ($P = 0.003$, $P = 0.001$).

Conclusion: The result of this study shows that there is a significant relation between stroke and periodontal index; however, there isn't any significant relation between stroke and gingival index.

Key Words: Gingivitis, periodontitis, risk factors, stroke

Received: April 2012
Accepted: June 2013

Address for correspondence:

Assistant Prof. Maryam
Alsadat Hashemipour,
Department of Oral
Medicine, Faculty of
Dentistry, Kerman University
of Medical Sciences.
Kerman, Iran
E-mail: m_hashemipour@
kmu.ac.ir

INTRODUCTION

Epidemiologic studies have shown that cerebrovascular accident (CVA) is the 3rd cause of death in developed countries with 0.8 % prevalence in whole population. Although, prevalence rate of stroke is reduced but yet is one of the most important health problem in United State.^[1,2]

CVA risk factors are hypertension, diabetes, smoking and cardiovascular disease. Not only urinary tract infections independently increase the risk of ischemic stroke, but also, mainly acute respiratory tract infection. Although, in many patients especially, young patients none of these risk factors doesn't exist. Today, inflammatory parameters and chronic and acute infectious diseases are considered as the risk factors for this disease.^[3,4] Though, there is not any single infection as a major independent predictor. In order to lower this burden it is essential to identify risk factors and respective preventive strategies besides the established stroke risk factors.^[5]

Periodontitis and gingivitis are two most infectious man diseases. Today, it is estimated that about 35%

Access this article online



Website: <http://drj.mui.ac.ir>

of people over 30-year-old in America and 50-55% of people in developing countries are involved with these diseases. These diseases are due to the practice of chronic bacterial infection and inflammatory response of host.^[3,4,6]

The relationship between periodontitis and CVA was first introduced by Mackenzi and Milard in 1963 but the importance of this issue was not understood.^[7] Periodontal infection can be a complex multiphase disease by associating this infection with organ systems such as cardiovascular, endocrine, reproductive, and respiratory systems.^[8]

Nowadays, periodontitis is considered as a risk factor for atherosclerosis and thromboembolic disease.^[6] Furthermore, periodontal infections may effect the onset and development of atherosclerosis by special mechanisms. Periodontitis and atherosclerosis have complex etiologic factors that are influenced by combination of hereditary and environmental factors. These diseases have many common risk factors and also have known similarity in essential pathogenic mechanism. Several studies have been carried out on dependence of periodontitis and stroke.^[9-14] Therefore, doing more investigation for determination of prevalence of periodontal disease in these patients for finding relationship between periodontal disease and CVA is completely necessary.^[7]

Research works in Iran have shown that there is a high prevalence rate of periodontal diseases in Iranian population, such that 1.2-10.5% of people have normal periodontium.^[15] Besides, the prevalence rate of stroke is between 0.43% and 0.74% in this region.^[16,17] Hence, regarding to the high prevalence of periodontal diseases in Iranian people and also noticing the fact that there is only one study about prevalence of periodontal disease and CVA patients in Iran,^[18] the present research work seems necessary to be carried out.

MATERIALS AND METHODS

A case-control study was conducted on 100 non-fatal stroke cases at the Shafa hospital, Kerman, Iran, and 100 hospitalized patients (control group). The matching between two groups was carried out according to this way that at first 100 patients from Neuroscience ward were examined by gingival and periodontal indexes. Then, based on the number of males and females and the age, 100 patients from others hospital wards were chosen.

Medical specialists diagnosed stroke by using brain imaging from magnetic resonance imaging and/or computerized tomography.^[18]

Using a University of North Carolina-15 manual probe, the clinical attachment level (CAL), the distance between the cemento-enamel junction (CEJ) and the probed base of the periodontal pocket, were recorded by a dentist. Data were collected by questionnaires, patient hospital records and clinical examinations. In the questionnaire, many items were included which are the demographic information (name, gender, profession, date of admission if hospitalized, diabetes mellitus, hypertension, heart valve disease, atrial fibrillation, peripheral arterial disease, previous stroke/ischemic attack, smoking, a positive family history of stroke, and renal failure) and information about cholesterol, triglyceride, Complete Blood Count (CBC) and evaluation of gingival and periodontal condition according to gingival index and periodontal disease index criteria.^[18]

Examination of all patients was carried out by a specially trained dentist for all subjects that had been trained 1 week in periodontal part of faculty. Examiner was tested in order to reduce the intra-examination error. Before the examination of each person, the goal of this study and method of examination was explained to patient and satisfaction form was gained from them. Furthermore, we gave confidence to patients that all of their information remain secret. All examination was carried out on patients with blue light, a mouth mirror and a periodontal probe (after isolation with cotton roles). For obvious reasons, the dentist could not be blinded for the patient's status. Smoking and nonsmoking and its rate was asked from patients and also blood pressure of patients and result of laboratory tests were gained from patient hospital records.^[18] The distance between the probed base of the pocket and the CEJ was considered as the main variable in CAL of periodontitis. In the following four sites of each available tooth, including distofacial papilla, facial gingival margin, mesio-facial papilla and lingual gingival margin, probing was performed and mean values were individually calculated. Attachment levels were analyzed as continuous variables. Using the Loe and Silness gingival index,^[19] gingivitis was determined by probing the teeth at four sites. For this purpose, scoring criterias for gingival are as follows:^[19] G0: normal gingival without inflammation, G1: mild inflammation, slight change in colour and texture and

no bleeding on probing, G2: moderate inflammation, shiny redness, edema and bleeding on very slight and light probing, G3: severe inflammation, marked shiny redness, edema, ulceration, and spontaneous bleeding on very slight and light probing.^[18,19] The data of this research were analyzed by use of multiple logistic regressions, Chi-square test, Fisher's test, *t*-test and SPSS11.5 software program. $P < 0.05$ was considered as significant.

RESULTS

In this study, 100 patient suffering from stroke as case group and 100 hospitalized patients as control group were selected. Average age of case group was 51.89 ± 15.51 and average of control group was 52.59 ± 17.3 . The case group included 42 (42%) males and 58 (58%) females, and in control group 44 males (44%) and 56 females (56%) were included. The mean time of examination after admission was 3.2 days for stroke patients and 2.9 days for

hospitalized control. Risk factors for cerebral ischemia are shown in Table 1.

In this investigation, the average of gingival index in men and women of case group in order was 1.22 ± 0.55 and 1.31 ± 0.55 with this fact that this number in women is more than men. However, there wasn't any significant difference between gender and gingival index ($P = 0.42$). Furthermore, the average of gingival index in case group was more than control group but there isn't any significant difference between them ($P = 0.64$) [Table 2].

In Table 3, the condition of gingival inflammation in two groups has been shown. In all criteria (mild, moderate and severe inflammation) in case group and in women is more than men, however, there isn't any significant relationship between gender and gingival index in two groups. Furthermore, there wasn't any significant difference about gingival inflammation between groups. According to our result, none of patients in two groups have not normal gingival.

Table 1: Risk factors for ischemic stroke in case and control groups

Ischemic stroke risk factors	Case group			Control group			P value
	Male (%)	Female (%)	Total (%)	Male (%)	Female (%)	Total (%)	
Familial history of ischemic stroke	12 (12)	12 (12)	24 (24)	3 (3)	2 (2)	5 (5)	0.001*
Diabetes mellitus	7 (7)	12 (12)	19 (19)	8 (8)	7 (7)	15 (15)	0.463
Peripheral cardiovascular disease	5 (5)	3 (3)	8 (8)	3 (3)	2 (2)	5 (5)	0.634
Hypertension	13 (13)	19 (19)	32 (32)	7 (7)	6 (6)	13 (13)	0.001*
Atrial fibrillation	7 (7)	3 (3)	10 (10)	2 (2)	5 (5)	7 (7)	0.453
Renal failure	6 (6)	4 (4)	10 (10)	1 (1)	1 (1)	2 (2)	0.004*
Smoking	15 (15)	0 (0)	15 (15)	9 (9)	1 (1)	10 (10)	0.213

*P value<0.05 (significant)

Table 2: Gingival index condition in case and control groups

Gingival index	Men (mean±SD)	Female (mean±SD)	Total (mean±SD)
Case group	1.22±0.51	1.31±0.55	1.27±0.53
Control group	1.1±0.32	1.11±0.21	1.1±0.26

Man Whitney, Degree of freedom: 2, P value: 0.64 (not significant)

Table 3: Gingival inflammation condition according to gingival index in case and control groups based on gender

Gingival index	Case group			Control group		
	Female	Male	Total	Female	Male	Total
Normal	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Mild inflammation	17 (29.3)	10 (23.8)	27 (27)	17 (29.3)	10 (23.8)	27 (27)
Moderate inflammation	34 (58.6)	27 (64.3)	61 (61)	31 (55.3)	26 (59.1)	57 (57)
Severe inflammation	7 (12.1)	5 (11.9)	12 (12)	5 (9)	3 (6.8)	8 (8)
Total	58 (58)	42 (52)	100 (100)	56 (56)	44 (44)	100 (100)

Chi2: 0.63, Degree of freedom: 2, P value: 0.72 (not significant)

Table 4, shows the frequency of gingivitis in case group according age groups. Relationship between age and gingivitis in case group was significant ($P = 0.007$).

Periodontal index in both groups in men was more than women and there was significant relationship between this index and gender ($P = 0.031$). Furthermore, periodontal index in case group was more than control group ($P = 0.001$) [Table 5]. The moderate and severe periodontitis in case group was more than control group and there was significant relationship between this index in two groups [Table 6]. Relationship between age and periodontal disease

in case group was significant ($P = 0.03$) [Table 7].

After considering age, gender, and other confounding factors in the multiple logistic regression models in the present work, it is found that periodontal index is associated with increased risk of stroke [Table 8].

We furthermore evaluated the association of the presence of periodontal disease with the risk factors of ischemic stroke. Results indicate that after omission of confounding factors only hypertension and renal failure are effective on periodontal and gingival condition in case group and significant association was observed ($P = 0.001$).

Table 4: Gingival inflammation condition in case group in different age groups

Age groups	Mild inflammation	Moderate inflammation	Severe inflammation	Total
Under 30 years	8 (66.6)	3 (25)	1 (8.4)	12
30-40 years	6 (35.29)	9 (52.94)	2 (11.76)	17
40-50 years	6 (30)	13 (65)	1 (5)	20
50-60 years	5 (20)	17 (68)	3 (12)	25
60-70 years	0	15 (83.33)	3 (16.67)	18
Upper 70 years	2 (25)	4 (50)	6 (25)	8
Total	27 (27.55)	61 (61.22)	12 (11.22)	100

Fisher's exact test: 24.35, Degree of freedom: 10, P value: 0.007 (significant)

Table 5: Periodontal index condition in case and control groups

Periodontal index	Male (mean±SD)	Female (mean±SD)	Total (mean±SD)
Case group	5.04±0.84	4.12±0.22	4.58±1.06
Control group	3.53±1.02	3.14±0.41	3.33±0.22

Man Whitney, Degree of freedom: 1, P value: 0.001 (significant)

Table 6: Frequency of periodontitis in case and control groups

Periodontal index	Case group			Control group			P value
	Female	Male	Total	Female	Male	Total	
Normal	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	—
Mild periodontitis	36 (62.1)	25 (59.2)	61 (61)	42 (61.90)	35 (66.67)	77 (27)	0.036*
Moderate periodontitis	19 (32.7)	16 (38.1)	35 (35)	14 (38.10)	9 (23.23)	23 (23)	0.003*
Severe periodontitis	3 (5.2)	1 (2.7)	4 (4)	0 (0)	0 (0)	0 (0)	0.001*
Total	58 (58)	42 (52)	100 (100)	56 (56)	44 (44)	100 (100)	—

* P value<0.05 (significant)

Table 7: Periodontitis condition in case group in different age groups

Age groups	Mild periodontitis	Moderate periodontitis	Severe periodontitis	Total
Under 30 years	10 (100)	0 (0)	0 (0)	10
30-40 years	12 (70.6)	4 (23.5)	1 (5.9)	17
40-50 years	12 (54.5)	8 (36.4)	2 (9.1)	22
50-60 years	15 (62.5)	8 (33.3)	1 (4.2)	24
60-70 years	8 (44.4)	10 (55.6)	0 (0)	18
Upper 70 years	3 (37.5)	5 (62.5)	0 (0)	8

χ^2 : 11.88, Degree of freedom:5, P value: 0.03 (significant)

Table 8: Gingival and periodontal index in case group in different age groups

Index	χ^2	df	P value	OR (CI)
Gingival	0.268	1	0.605	0.8 (0.356-1.82)
Periodontal	5.245	1	0.032*	2.7 (1.26-6.4)

*P value<0.05 (significant), df: Degree of freedom, OR: odds ratio; CI: confidence interval; Multiple logistic regression

DISCUSSION

Chronic infections have been reported to be as risk factors for ischemic stroke. However, the association of ischemic stroke with periodontal disease, which is also a chronic inflammatory disease, needs to be evaluated.^[18] This study showed that prevalence of gingival index and periodontal index in case group is more than control group.

In 2000, the American general surgeon committee published the first report about oral health and hygiene with emphasize on this point that oral hygiene is beyond having intact teeth and is the main part of general health. Based on surgeon idea, mouth is the way for entering microbial infection that influence on general health situation. Furthermore, mouth is a vital and functional tissue that can have important role focus for general health, and having healthy life.^[20]

During two recent decades, studies about oral hygiene, atherosclerosis and CVA have been carried out.^[9-18,21-25] Stroke risk factors include aging, smoking, race, hypertension, diabetes mellitus, cardiovascular disease, alcoholism, inflammatory disease, and Obesity.^[11] Inflammatory risk factors of stroke have been assessed and in the opinion of researchers periodontitis is dependent to stroke.^[14,26]

There are several potential mechanisms for relationship between periodontal disease and stroke.^[7,10,26,27] Periodontitis causes an increase in the systemic inflammation that leads to atherosclerosis and cerebrovascular attacks. Acute periodontal inflammation may cause a stage of prothrombin, bacteremia, platelet activation, increasing in coagulator's factors and risk of stroke. Furthermore, periodontal microorganism was observed in atheroma. The endotoxin of this microorganisms can destroy endothelial cells and cause smooth muscle proliferation.^[11,26] Periodontal disease produces inflammatory markers and coagulator's factors such as C-reactive protein, fibrinogen and increases platelets aggregation and cause atherosclerosis

and thrombosis.^[11,28] Furthermore, chronic gingival infections decreases glucose tolerances and decreases blood lipids.^[18]

The periodontal infection have a main role by creation stable bacterial environment and take part in inflammatory processes derived from monocytes and macrophages that cause atheromatosis and narrowing of blood vessels. Furthermore, bacteremia deriving from subgingival and supragingival plaque can increase platelets aggregation by taking part in thromboses formation and subsequently embolism which are the main causes of stroke.^[29]

Because of multifactorial nature of dental infection and CVA, founding a distinguished factor is difficult and results have contradiction with each other. The most principle problem of these studies is insufficient control on numerous confounding factors that cause lower estimation of risk factors.

In this study, periodontitis was indicated as an independent risk factor for stroke, especially, in men and young subjects. Associations between cerebral ischemia and periodontitis was distinguished by several epidemiological studies.^[9,22-24]

One meta-analysis of retrospective and prospective study has shown that periodontal disease may increase risk of CVA up to 20%. Similarly, reported relative risk between periodontal disease and stroke from 2.85 to 1.74 is changeable.^[30]

The study by Mendez *et al.* on 80 subjects during 25-30 years follow-up showed that patients with clinically apparent periodontal disease, are 2.27 times more than others in the risk of peripheral vessels disease.^[31] These researchers concluded that periodontal disease is an independent risk factor for peripheral vessels disease. Furthermore, in a report by Wu *et al.* there was clear relationship between periodontal disease and stroke.^[32] In other retrospective studies by Morrison *et al.* an indistinctive relation between periodontal disease and stroke has been shown. This evaluation detected that this relationship increases to 2.12% for fatal ischemia.^[25]

It was concluded by Janket *et al.* that there is a small but significant increase in the risk of cardiovascular disease among patients with periodontal disease at baseline.^[33] In another research by Grau *et al.* periodontitis were distinguished as an independent risk factor for cerebral ischemia, especially, in men and young patients.^[9] This finding is similar to that

obtained by Kim *et al.*, Pradeep *et al.*, Dörfer *et al.*, Geerts *et al.*, which are in agreement with the present study.^[21,22,34,35]

Jimenez *et al.* in a current study revealed an association between history of periodontitis but not current periodontal inflammation and incidence of cerebrovascular disease in men, independent of established cardiovascular risk factors, particularly among men aged <65 years.^[36]

One study has reported that men with periodontal disease in comparison with men without periodontal disease almost 3 times more likely to have stroke risk.^[7] Furthermore, study of Joshipura *et al.* on 41380 men without cardiovascular disease during 12 years follow-up showed that during this period, 349 persons were affected with stroke. Men with gingival index more than 1.5 were under the stroke risk about 1.3 times more than others.^[13]

Abolfazli *et al.* showed that patients (case group) have higher loss of attachment than controls, whereas gingivitis isn't associated with the risk of cerebral ischemia. Furthermore, the more severe periodontitis was detected among men in patients with cerebral ischemia.^[18]

In similar studies,^[9,10,13,18] gingivitis and periodontitis were reported in men more than women. That is because of less oral hygiene compliance in men and infrequent referring to dentist.

Present study shows that there isn't significant relationship between gingival indexes between 2 groups. Although, in some study severe gingivitis have been propounded as a more important risk factor in CVA patient in comparison to periodontitis.^[9,11] Researchers believe that gingivitis is a start to periodontal diseases, although, in one cohort study, any association between these two factors was not found.^[32]

In this study, in fully adjusted logistic regression analysis, a probing pocket depth of >4.5 mm was found to be the most significant factor for stroke (odds ratio = 2.7; confidence interval = 1.26-6.4; $P = .032$).

Sfyroeras *et al.* carried out six prospective and seven retrospective studies about association between periodontal disease and stroke. These researchers showed that overall adjusted risk of stroke in subjects with periodontitis is 1.47 times higher than in subjects without periodontitis in prospective studies (95% confidence interval, 1.13-1.92; $P = 0.0035$)

and 2.63 times (95% confidence interval, 1.59-4.33; $P = 0.0002$) in retrospective studies.^[37]

Studies that assessed tooth loss or periodontal disease as risk factor of stroke lead to different results and the relation between oral condition and CVA is yet unclear. In the absence of any cause-effect relation, this relationship may explain by common risk factors. Therefore, more epidemiologic studies are needed to discover the role of nutrition and other inflammatory mediators.

CONCLUSION

The result of this study showed that there is significant relation between stroke and periodontal index, however, there isn't any significant relation between stroke and gingival index. Furthermore, it was found that periodontal index in women and age groups more 40 years are high.

ACKNOWLEDGMENTS

This study was supported by the Kerman Neuroscience Research Center. The authors would like to thank the Research Deputy for their financial support.

REFERENCES

1. Greenburg M, Glick M. *Burket's Oral Medicine*. 11th ed. Hamilton: BC Decker; 2009. p.98, 115-6.
2. Little JW, Falace DA, Miller CS, Rhodus NL. *Dental Management of the Medically Compromised Patients*. 2nd ed. Philadelphia: WB Saunders; 2003. p. 121-48.
3. Lindsberg PJ, Grau AJ. Inflammation and infections as risk factors for ischemic stroke. *Stroke* 2003;34:2518-32.
4. Chala S, Abouqal R, Abdallaoui F. Prevalence of apical periodontitis and factors associated with the periradicular status. *Acta Odontol Scand* 2011;69:355-9.
5. Elkind MS, Ramakrishnan P, Moon YP, Boden-Albala B, Liu KM, Spitalnik SL, *et al.* Infectious burden and risk of stroke: The northern Manhattan study. *Arch Neurol* 2010;67:33-8.
6. Scannapieco FA. Position paper of The American Academy of Periodontology: Periodontal disease as a potential risk factor for systemic diseases. *J Periodontol* 1998;69:841-50.
7. Beck J, Garcia R, Heiss G, Vokonas PS, Offenbacher S. Periodontal disease and cardiovascular disease. *J Periodontol* 1996;67:1123-37.
8. Saini R, Marawar PP, Shete S, Saini S. Periodontitis, a true infection. *J Glob Infect Dis* 2009;1:149-50.
9. Grau AJ, Buggle F, Ziegler C, Schwarz W, Meuser J, Tasman AJ, *et al.* Association between acute cerebrovascular ischemia and chronic and recurrent infection. *Stroke* 1997;28:1724-9.
10. Howell TH, Ridker PM, Ajani UA, Hennekens CH, Christen WG. Periodontal disease and risk of subsequent cardiovascular disease in U.S. male physicians. *J Am Coll Cardiol* 2001;37:445-50.

11. Desvarieux M, Schwahn C, Völzke H, Demmer RT, Lüdemann J, Kessler C, *et al.* Gender differences in the relationship between periodontal disease, tooth loss, and atherosclerosis. *Stroke* 2004;35:2029-35.
12. Grau AJ, Becher H, Ziegler CM, Lichy C, Buggle F, Kaiser C, *et al.* Periodontal disease as a risk factor for ischemic stroke. *Stroke* 2004;35:496-501.
13. Josphipura KJ, Hung HC, Rimm EB, Willett WC, Ascherio A. Periodontal disease, tooth loss, and incidence of ischemic stroke. *Stroke* 2003;34:47-52.
14. Syrjänen J, Peltola J, Valtonen V, Iivanainen M, Kaste M, Huttunen JK. Dental infections in association with cerebral infarction in young and middle-aged men. *J Intern Med* 1989;225:179-84.
15. Arabi SR, Torkzaban P, Gholami L, Taghavi M, Hatami M. Comparative evaluation of periodontal indices in patients with ischemic heart disease and positive myocardial perfusion scan. *Dent J Hamadan Univ* 2010;1:17-20.
16. Oveisgharan S, Sarrafzadegan N, Shirani S, Hosseini S, Hasanzadeh P, Khosravi A. Stroke in Isfahan, Iran: Hospital admission and 28-day case fatality rate. *Cerebrovasc Dis* 2007;24:495-9.
17. Ghandehari K, Moud ZI. Incidence and etiology of ischemic stroke in Persian young adults. *Acta Neurol Scand* 2006;113:121-4.
18. Abolfazli N, Ghoreishizadeh A, Ayramlu H, Ghavimi M, Ghoreishizadeh M, Salehsaber F. Periodontal disease and risk of cerebral ischemic stroke. *J Neurol Sci* 2011;28:307-16.
19. Newman MG, Takei HH, Carranza FA. Carranza's Clinical Periodontology. 10th ed. Hamilton WB Saunders; 2006. p. 112-22, 333-53.
20. Rose LF, Mealey B, Minsk L, Cohen DW. Oral care for patients with cardiovascular disease and stroke. *J Am Dent Assoc* 2002;133:37S-44.
21. Pradeep AR, Hadge P, Arjun Raju P, Shetty SR, Shareef K, Guruprasad CN. Periodontitis as a risk factor for cerebrovascular accident: A case-control study in the Indian population. *J Periodontol Res* 2010;45:223-8.
22. Dörfer CE, Becher H, Ziegler CM, Kaiser C, Lutz R, Jörss D, *et al.* The association of gingivitis and periodontitis with ischemic stroke. *J Clin Periodontol* 2004;31:396-401.
23. Khader YS, Albashaireh ZS, Alomari MA. Periodontal diseases and the risk of coronary heart and cerebrovascular diseases: A meta-analysis. *J Periodontol* 2004;75:1046-53.
24. Lee HJ, Garcia RI, Janket SJ, Jones JA, Mascarenhas AK, Scott TE, *et al.* The association between cumulative periodontal disease and stroke history in older adults. *J Periodontol* 2006;77:1744-54.
25. Morrison HI, Ellison LF, Taylor GW. Periodontal disease and risk of fatal coronary heart and cerebrovascular diseases. *J Cardiovasc Risk* 1999;6:7-11.
26. Elter JR, Offenbacher S, Toole JF, Beck JD. Relationship of periodontal disease and edentulism to stroke/TIA. *J Dent Res* 2003;82:998-1001.
27. Ziegler CM, Schwarz W, Grau A, Buggle F, Hassfeld S, Mühling J. Odontogenic focus as the etiology of cerebral ischemia. *Mund Kiefer Gesichtschir* 1998;2:316-9.
28. Noack B, Genco RJ, Trevisan M, Grossi S, Zambon JJ, De Nardin E. Periodontal infections contribute to elevated systemic C-reactive protein level. *J Periodontol* 2001;72:1221-7.
29. Mealey BL. Influence of periodontal infections on systemic health. *Periodontol* 2000 1999;21:197-209.
30. Meurman JH, Sanz M, Janket SJ. Oral health, atherosclerosis, and cardiovascular disease. *Crit Rev Oral Biol Med* 2004;15:403-13.
31. Mendez MV, Scott T, LaMorte W, Vokonas P, Menzoian JO, Garcia R. An association between periodontal disease and peripheral vascular disease. *Am J Surg* 1998;176:153-7.
32. Wu T, Trevisan M, Genco RJ, Dorn JP, Falkner KL, Sempos CT. Periodontal disease and risk of cerebrovascular disease: The first national health and nutrition examination survey and its follow-up study. *Arch Intern Med* 2000;160:2749-55.
33. Janket SJ, Baird AE, Chuang SK, Jones JA. Meta-analysis of periodontal disease and risk of coronary heart disease and stroke. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;95:559-69.
34. Kim HD, Sim SJ, Moon JY, Hong YC, Han DH. Association between periodontitis and hemorrhagic stroke among Koreans: A case-control study. *J Periodontol* 2010;81:658-65.
35. Geerts SO, Legrand V, Charpentier J, Albert A, Rompen EH. Further evidence of the association between periodontal conditions and coronary artery disease. *J Periodontol* 2004;75:1274-80.
36. Jimenez M, Krall EA, Garcia RI, Vokonas PS, Dietrich T. Periodontitis and incidence of cerebrovascular disease in men. *Ann Neurol* 2009;66:505-12.
37. Sfyroeras GS, Roussas N, Saleptsis VG, Argyriou C, Giannoukas AD. Association between periodontal disease and stroke. *J Vasc Surg* 2012;55:1178-84.

How to cite this article: Hashemipour MA, Afshar AJ, Borna R, Seddighi B, Motamedi A. Gingivitis and periodontitis as a risk factor for stroke: A case-control study in the Iranian population. *Dent Res J* 2013;10:613-9.

Source of Support: Nil. **Conflict of Interest:** None declared.