

Original Article

Comparative evaluation of bispectral index system after sedation with midazolam and propofol combined with remifentanyl versus ketamine in uncooperative during dental procedures

Alireza Eshghi¹, Mehrnaz Mohammadpour², Nasser Kaviani³, Dana Tahririan⁴, Najmeh Akhlaghi¹

¹Torabinejad Dental Research Center and Departments of Pediatric Dentistry, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, ²Department of Pediatrics, School of Dentistry, Shahr-e-kord University of Medical Sciences, Shahr-e-kord, ³Torabinejad Dental Research Center and Oral and Maxillofacial Surgery, School of Dentistry, Isfahan University of Medical Sciences, ⁴Dental Materials Research Center and Departments of Pediatric Dentistry, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

ABSTRACT

Background: Proper analgesic agents should be used in combination with sedative agents. Remifentanyl is a synthetic narcotic/analgesic agent with a short duration effect and decreases the risk of apnea during recovery. Bispectral index system (BIS) is a new noninvasive technique for the evaluation of the depth of sedation. The aim of present clinical trial was to evaluate and compare the efficacy of intravenous sedation with propofol/midazolam/remifentanyl (PMR) in comparison to propofol/midazolam/ketamine (PMK) for dental procedures in children 3-7 years of age.

Materials and Methods: In this clinical trial, 32 healthy uncooperative children who were candidates for dental treatments under sedation were randomly divided into two groups. Intravenous sedation was induced with PMR in one group and with PMK in the other group. After injection and during procedure BIS index, heart rate and respiratory rate, blood pressure, and oxygen saturation was evaluated every 5 min. After the procedure, recovery time was measured. Data were analyzed with ANOVA, Friedman, Wilcoxon, and *t*-test.

Results: The BIS value was significantly low in ketamin group ($P = 0.003$) but respiratory rates and heart rates were same in both groups with no statistical difference ($P = 0.884$, $P = 0.775$). The recovery time was significantly shorter in remifentanyl group ($P = 0.008$ and $P = 0.003$).

Conclusion: It can be concluded that intravenous sedation technique with PMR combination induces effective and safe sedation, with less pain and more forgetfulness and a shorter recovery time for children 3-7 years of age during dental procedures.

Key Words: Bispectral index, ketamine, midazolam, propofol, remifentanyl

Received: April 2015
Accepted: September 2015

Address for correspondence:
Dr. Mehrnaz Mohammadpour,
Department of Pediatrics,
School of Dentistry,
Shahr-e-kord University
of Medical Sciences,
Shahr-e-kord, Iran.
E-mail: mehrnaz_m_d@
yahoo.com

INTRODUCTION

Intravenous sedation is the second most comfortable, effective, and the safest technique after inhalation sedation; however, the technique should be used only by trained personnel.^[1] Sedative techniques do not

involve intubation, contrasting general anesthesia and the recovery period is short.^[2] There are basically two techniques available for intravenous sedation: Use of

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Eshghi A, Mohammadpour M, Kaviani N, Tahririan D, Akhlaghi N. Comparative evaluation of bispectral index system after sedation with midazolam and propofol combined with remifentanyl versus ketamine in uncooperative during dental procedures. Dent Res J 2016;13:1-6.

Access this article online



Website: www.drj.ir
www.drjjournal.net
www.ncbi.nlm.nih.gov/pmc/journals/1480

only one medication, which is usually benzodiazepine and a combination of several medicines.^[1] In sedation, it is of the utmost importance to achieve analgesia. Due to the limitations of the use of local anesthetic agents such as overdose and reaching the toxic levels of medications, proper analgesic agents should be used in combination with sedative agents.^[3]

The most commonly medications used in sedation are ketamine, midazolam, and propofol.^[4-6] Ketamine has an analgesic effect but may cause complications during recovery, including severe listlessness, nausea, delirium, nystagmus, and severe muscle spasms. It is contraindicated in patients with a history of convulsions.^[5,6] In contrast, propofol does not have an analgesic effect but may increase the risk of respiratory depression, body irritation, crying, and coughing during the procedure and anxiety during recovery, with no nausea.^[6,7] Midazolam has no analgesic effect but may induce forgetfulness after the procedure.^[7] Remifentanyl is a synthetic narcotic/analgesic agent with a strong analgesic effect which decreases the risk of apnea during recovery; however, it has a short duration of effect.^[8] Since it results in minor changes in cardiac waves and leads to rapid recovery, remifentanyl is a good choice for cardiac and debilitated patients.^[9,10]

Bispectral index system (BIS) is a new noninvasive technique for the evaluation of the depth of sedation and may be a proper technique for evaluation of children undergoing a sedative technique.^[11,12] The electroencephalography (EEG) device reports the waves numerically in a range of 0-100, in which 100 indicates full consciousness and a value between 60 and 90 indicates adequate sedation.^[13,14] These values should be in the range of 40-60 in general anesthesia, 60-70 in profound sedation, and 70-90 in moderate sedation.^[14] There are only a few studies available on intravenous sedation of children for dental procedures. Combining medications results in the use of lower doses and the risks associated with medications decrease.^[5-8,15] A combination of propofol, fentanyl and midazolam resulted in more effective sedation, with shorter recovery, compared to the use of propofol alone^[15] and the use of a combination of propofol and remifentanyl was safe, effective, and acceptable.^[8] Regarding the , inconsistent results of propofol^[5-8,12]. Remifentanyl is a new medication and its combination with midazolam increases forgetfulness after the procedure compared to midazolam alone.^[16] Further studies are needed to determine the best combination

of drugs for sedation in children.^[5-8,15,16] The present study was undertaken to evaluate and compare the efficacy of a combination of propofol/midazolam/ketamine (PMK) with that of a combination of propofol/midazolam/remifentanyl (PMR) in sedating children during dental procedures by means of BIS technique.

MATERIALS AND METHODS

Participants

In the present prospective, double-blind clinical trial a total of 32 uncooperative children (one or two negatives based on Frankel behavioral rating scale^[1]), aged 3-7, who referred to Department of Hospital Dentistry in Isfahan Dental School selected randomly. The inclusion criteria were healthy children who had no specific systemic disease and were in the category I of the American Society of Anesthesiology. At the time of sedation, children did not have a common cold or any airway problems. Patients with extraction or who needed dental work time more than 45 min were excluded. The sample size in each group was 16 subjects at a significance level of 0.05 and a power of 80% ($\alpha = 0.05$, $\beta = 0.20$), using the below formula. This was estimated to show a six difference in the mean of BIS index between the two groups.^[10]

$$n = \frac{(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta})(\delta_1^2 + \delta_2^2)}{d^2} \quad \delta_1 = \delta_2 = \frac{0}{5} \quad d = 6$$

Sedation protocols and patient monitoring

After obtaining informed consent from the parents, they received the necessary instructions for the sedative procedures. All the patients were asked to refer in the morning, in a fasting state (at least 5 h not to eat by mouth or non per os), for the dental procedures which performed at Department of Hospital Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran.

Randomization was carried out as follows: On the day of the procedure each subject was given a code of which only the anesthesiologist was aware. Neither subjects and nor other researchers were aware of codes. Patients were randomly divided into two groups based on the odd or even codes. In the group with even codes, propofol/midazolam/ketamine was administered intravenously in the following doses: Ketamine (0.5 mg/kg); propofol (0.5 mg/kg); and midazolam (0.01 mg/kg).

In the group with odd codes, midazolam/propofol/remifentanyl was administered intravenously in

the following doses: Midazolam (0.01 mg/kg); propofol (0.5 mg/kg); and remifentanil were pumped intravenously at (0.1 µg/kg/min).

The depth of sedation, BIS, heart rate, the number of respirations, blood pressure, and oxygen saturation percentages were recorded every 5 min by the anesthesiologist. Therefore, a complete monitoring which is essential for general anesthesia or deep sedation was performed for both groups. The conditions for carrying out the procedure were evaluated by another researcher based on the Dental Sedation Teachers Group (DSTG)^[17,18] scale and recorded in special data sheets. At the end of the procedure, each patient was monitored in the recovery room and the time needed for recovery was recorded. After achieving the necessary conditions for being discharged based on the postanesthetic discharge scoring system,^[1] each patient was discharged based on thorough counseling and provision of a phone number for emergency calls.

Statistical methods

After collection of data the codes were deciphered and data were analyzed with SPSS (version 20, Chicago, IL, USA), using Friedman's and Wilcoxon's tests, ANOVA, *t*-test and Pearson's correlation coefficient. The level of the confidence interval was considered at 95%.

Ethical aspect

The protocol and consent form were approved by Research Committee of Isfahan University of Medical Science (code number 393398) and Registered in Iranian Registry of Clinical Trials (registration code #IRCT2015042512848N2). It should be noted that drug doses were completely based on medical literature, and careful patient care was provided during the work. Moreover, attempts to treat a child

in ordinary dental appointment were not successful due to lack of cooperation, parents informed of the treatment under sedation, and written consent was obtained. Patients pay only the cost of dental treatments, and no additional charges would be incurred for this study.

RESULTS

Study population and mean total dose of drugs

A total of 32 children aged 3-7 years, consisting of 17 girls (52%) and 15 boys (48%), were included in the present study. The mean age of the subjects was 4.36 ± 1.6 years. *t*-test showed no significant differences in gender, duration of the procedures and body weights between the two groups [Table 1].

The bispectral index system values between groups

The mean BIS index in the remifentanil group 5 min after the initiation of sedation was 68.62 ± 10.24 , which first decreased during the sedation time but began to increase approximately 20 min after the initiation of sedation, finally reaching 69.71 ± 4.57 . In the ketamine group, the mean BIS index 5 min after initiation of sedation was 50.08 ± 8.39 , which first decreased in a manner similar to that in the remifentanil group but increased with time and finally reached a value close to that at the initiation time. Analyses of covariance for the repeated data showed significant differences in BIS between the two groups ($P = 0.003$), with significantly higher BIS index values in the remifentanil group compared to the ketamine group during the whole procedures [Table 2].

The bispectral index system values intra-groups in different time of procedure

The BIS index values were compared at nine different time intervals separately in each group, demonstrating no significant differences between the remifentanil and ketamine groups (intra-group) in different time intervals ($P = 0.505$ and $P = 0.577$, respectively). In all the subjects in both groups the DSTG sedation score^[18] at all the nine different time intervals was 5 (eyes closed, no response to mild physical stimuli) and therefore, the conditions for carrying out the

Table 1: Demographics and the mean working time

Group	Mean age	Male	Female	Mean working time (min)
Remifentanil	1/20±3/72	7	9	8/53±39/62
Ketamin	1/75±5/50	8	8	9/73±40/83
<i>P_v</i>	0/037		0/9	0/742

Table 2: The BIS index values at different time intervals of the study (in min) in the two groups

Group	Interval									<i>P</i>
	5	10	15	20	25	30	35	40	45	
Remifentanil	68.62±10.24	67.62±9.21	56.54±8.48	65.15±6.37	65.31±6.72	65.75±4.28	68±5.17	67.13±5.30	69.71±4.57	0.505
Ketamine	50.08±8.39	53.58±8.45	51.58±11.09	49.25±9.27	49.82±10.71	49.09±8.74	48.33±6.36	50.33±9.23	50±11.95	0.577

BIS: Bispectral index system.

procedures were deemed “good” in all the subjects (the patient fully cooperative and proper sedation).

Safety profile

The heart rates in the two groups were evaluated and compared with repeated-measures ANOVA, revealing no significant differences between the two groups ($P = 0.884$).

The respiratory rates were evaluated at nine different time intervals using repeated-measures ANOVA, and no significant differences were found between the two groups ($P = 0.775$).

In all the subjects in both groups, the oxygen saturation percentage (PO_2 saturation) at different time intervals was 97-100%, and in none of the subjects it decreased to levels below 97%.

Procedure-related times

The mean recovery times from the time of medication administration in the remifentanyl and ketamine groups were 9.23 ± 2.77 and 30.83 ± 5.96 min, respectively. *t*-test revealed a significant difference between the two groups ($P < 0.001$). Pearson’s correlation coefficient did not reveal any significant correlation between the duration of the procedure and the recovery time ($r = 0.255$, $P = 0.218$).

Satisfaction survey

No psychomimetic and other complications, including severe sleepiness, negative reactions, and respiratory depression, were observed while the child was in the dental operatory or in the recovery room. Only a few cases of severe nausea and vomiting were reported in a few subjects in the remifentanyl group, which were resolved by prescribing plasil (metoclopramide).

DISCUSSION

In the present blind clinical trial, the PMR combination was significantly more effective ($P = 0.003$) and was associated with less recovery time than PMK ($P < 0.001$). To our knowledge, there are a limited number of studies have compared the two drugs, remifentanyl and ketamine (alone or in combination with other drugs). Several studies have shown that incorporation of remifentanyl shortens the recovery time, consistent with the results of the present study.^[15,19-22]

However, Moerman concluded that the addition of remifentanyl to propofol during spontaneous

ventilation had no benefits compared with the use of propofol alone. Although the addition of remifentanyl resulted in a dose reduction of propofol, using propofol alone accompany with less recovery time and better patient satisfaction.^[9]

In the present study, there were no significant differences in heart rates and respiratory rates between the two groups. Several studies have been conducted about side effects of remifentanyl or other opioids. However, there are many discrepancies in results. For example, one study reported that the side effects of remifentanyl were not more than nonopioid drugs.^[23] While others demonstrated that complications such as breathing and conscious reduction were observed in the use of remifentanyl.^[8,9] The combination of remifentanyl with midazolam increases forgetfulness after the procedure compared to midazolam alone.^[16] It seems that PMR combination had more satisfactory results than propofol/remifentanyl.

BIS scores in the PMK group were lower at all the time intervals than those in the PMR group, and more profound sedation had been achieved in the PMK group. The results of the present study were somewhat close to other studies such as Kramer *et al.* who compared the sedative effect of the combinations of remifentanyl/propofol and ketamine/propofol. They found that both groups had the same respiratory and cardiac complications; however, the duration of sedative effect and the time of recovery for the ketamine group were significantly higher than remifentanyl.^[24] Moreover, Berkenbosch *et al.* reported that the combination of remifentanyl and propofol had a well sedative effect and fast recovery with no serious side effects.^[25] The most commonly technique to evaluate the depth of sedation is to assess clinical signs such as patient movement and response to verbal and physical stimuli which principally depend on the patient’s personal reports and are considered an indirect indicator of the effect of medications on the brain.^[1] Although EEG is one of the most straightforward techniques to evaluate brain activity during sedation, this technique is difficult and costly. BIS is a new noninvasive technique for the evaluation of the depth of sedation and might be a proper technique for evaluation of children undergoing a sedative technique.^[11,12] Studies have shown that a BIS score ≥ 60 indicates an appropriate depth of sedation, with no possibility of accidental awakening during the procedure,^[26] which was also shown in the present study. In the remifentanyl group, the subjects were in

a profound sedation state at different time intervals; however, in the ketamine groups, the subjects were in general anesthesia state at all the intervals. Such a situation might be explained by the fact that the individual physiologic responses of children to a fixed dose are different. In other words, despite the use of a fixed dose for sedation, some children go into general anesthesia and the results of the present study showed that such a possibility is definitely higher with ketamine compared to remifentanyl. Therefore, ketamine and remifentanyl should be used for general anesthesia and deep sedation in the presence of proper tools under monitoring. Since there were no side effects, such as respiratory muscles spasms, apnea during recovery and laryngeal ulcers due to the absence of intubation, in the sedation technique compared to general anesthesia technique, more extensive a comprehensive studies are suggested to evaluate intravenous sedation techniques for dental procedures. The administration of PMK for general anesthesia and PMR for sedation purposes could be suggested.

CONCLUSION

It can be concluded that intravenous sedation technique with a combination of midazolam, propofol, and remifentanyl induces effective and safe sedation, with less pain and more forgetfulness and a less recovery time for children 3-7 years of age during dental procedures.

ACKNOWLEDGMENTS

The authors would like acknowledge to the staff at the Department of Hospital Dentistry in School of Dentistry, Isfahan University of Medical Sciences. This study was based on a thesis to be submitted to Dental School, in partial fulfillment of the requirements of Specialist in Pediatric Dentistry. This study was funded by Vice Chancellery for Research, (Grant number 393398, IUMS).

Financial support and sponsorship

Torabinejad Dental Research Center, Isfahan University of Medical Sciences.

Conflicts of interest

No one of those who were involved in conception, designing, planning, running trial, interpretation, drafting manuscript and submission are funded or supported by any manufacturer or any other related affairs organization.

REFERENCES

1. Golpayegani MV, Dehghan F, Ansari G, Shayeghi S. Comparison of oral midazolam-ketamine and midazolam-promethazine as sedative agents in pediatric dentistry. *Dent Res J (Isfahan)* 2012;9:36-40.
2. American Dental Association. Guidelines for Teaching Pain Control and Sedation to Dentists and Dental Students. Chicago: ADA; 2007.
3. Toomarian L, Salem K, Ansari G. Assessing the sedative effect of oral vs submucosal meperidine in pediatric dental patients. *Dent Res J (Isfahan)* 2013;10:173-9.
4. Heard C, Smith J, Creighton P, Joshi P, Feldman D, Lerman J. A comparison of four sedation techniques for pediatric dental surgery. *Paediatr Anaesth* 2010;20:924-30.
5. Shah A, Mosdossy G, McLeod S, Lehnhardt K, Peddle M, Rieder M. A blinded, randomized controlled trial to evaluate ketamine/propofol versus ketamine alone for procedural sedation in children. *Ann Emerg Med* 2011;57:425-33.e2.
6. Godambe SA, Elliot V, Matheny D, Pershad J. Comparison of propofol/fentanyl versus ketamine/midazolam for brief orthopedic procedural sedation in a pediatric emergency department. *Pediatrics* 2003;112(1 Pt 1):116-23.
7. Rai K, Hegde AM, Goel K. Sedation in uncooperative children undergoing dental procedures: A comparative evaluation of midazolam, propofol and ketamine. *J Clin Pediatr Dent* 2007;32:1-4.
8. Abu-Shahwan I, Mack D. Propofol and remifentanyl for deep sedation in children undergoing gastrointestinal endoscopy. *Paediatr Anaesth* 2007;17:460-3.
9. Moerman AT, Struys MM, Vereecke HE, Herregods LL, De Vos MM, Mortier EP. Remifentanyl used to supplement propofol does not improve quality of sedation during spontaneous respiration. *J Clin Anesth* 2004;16:237-43.
10. Ganzberg S, Pape RA, Beck FM. Remifentanyl for use during conscious sedation in outpatient oral surgery. *J Oral Maxillofac Surg* 2002;60:244-50.
11. Overly FL, Wright RO, Connor FA, Jay GD, Linakis JG. Bispectral analysis during deep sedation of pediatric oral surgery patients. *J Oral Maxillofac Surg* 2005;63:215-9.
12. Powers KS, Nazarian EB, Tapyrik SA, Kohli SM, Yin H, van der Jagt EW, *et al.* Bispectral index as a guide for titration of propofol during procedural sedation among children. *Pediatrics* 2005;115:1666-74.
13. Johansen JW, Sebel PS. Development and clinical application of electroencephalographic bispectrum monitoring. *Anesthesiology* 2000;93:1336-44.
14. Rosow C, Manberg PJ. Bispectral index monitoring. *Anesthesiol Clin North America* 2001;19:947-66, xi.
15. VanNatta ME, Rex DK. Propofol alone titrated to deep sedation versus propofol in combination with opioids and/or benzodiazepines and titrated to moderate sedation for colonoscopy. *Am J Gastroenterol* 2006;101:2209-17.
16. Garip H, Gürkan Y, Toker K, Göker K. A comparison of midazolam and midazolam with remifentanyl for patient-controlled sedation during operations on third molars. *Br J Oral Maxillofac Surg* 2007;45:212-6.

17. Kaviani N, Ashrafi S, Jabbarifar SE, Ghaffari E. The Efficacy of Two Intravenous Sedative Drugs in Management of Uncooperative Children for Dental Treatments. *J Dent (Shiraz)* 2015;16 1 Suppl:29-34.
18. Robinson E, Kelly G. DSTG symposium report 14th May 2013, Cardiff millennium stadium. *SAAD Dig* 2014;30:51-4.
19. Hong MJ, Sung IK, Lee SP, Cheon BK, Kang H, Kim TY. Randomized comparison of recovery time after use of remifentanyl alone versus midazolam and meperidine for colonoscopy anesthesia. *Dig Endosc* 2015;27:113-20.
20. Rudner R, Jalowiecki P, Kawecki P, Gonciarz M, Mularczyk A, Petelencz M. Conscious analgesia/sedation with remifentanyl and propofol versus total intravenous anesthesia with fentanyl, midazolam, and propofol for outpatient colonoscopy. *Gastrointest Endosc* 2003;57:657-63.
21. Collini S, Pinto G, Lejeune L, Di Carlo S, Meloncelli S, Barraco G, *et al.* Neurosedation in dentistry of the disabled patient: The use of midazolam, propofol, and remifentanyl. *Minerva Stomatol* 2006;55:99-113.
22. Kramer KJ, Ganzberg S, Prior S, Rashid RG. Comparison of propofol-remifentanyl versus propofol-ketamine deep sedation for third molar surgery. *Anesth Prog* 2012;59:107-17.
23. Pinsker MC, Carroll NV. Quality of emergence from anesthesia and incidence of vomiting with remifentanyl in a pediatric population. *Anesth Analg* 1999;89:71-4.
24. McDermott NB, VanSickle T, Motas D, Friesen RH. Validation of the bispectral index monitor during conscious and deep sedation in children. *Anesth Analg* 2003;97:39-43.
25. Berkenbosch JW, Graff GR, Stark JM, Ner Z, Tobias JD. Use of a remifentanyl-propofol mixture for pediatric flexible fiberoptic bronchoscopy sedation. *Paediatr Anaesth* 2004;14:941-6.
26. Tschiedel E, Müller O, Schara U, Felderhoff-Müser U, Dohna-Schwake C. Sedation monitoring during open muscle biopsy in children by Comfort Score and Bispectral Index — A prospective analysis. *Paediatr Anaesth* 2015;25:265-71.