

Original Article

The association between periodontal disease parameters and severity of atherosclerosis

Mohammad Ketabi^{1,2}, Fatemeh Rashidi Meybodi³, Mohammad Reza Asgari⁴

Departments of ¹Periodontics and ³Oral Diseases, Faculty of Dentistry, Isalmic Azad University, Isfahan (Khorasgan) Branch, Isfahan, Iran, ²Visiting Professor, Department of Periodontology, Faculty of Dentistry University of Toronto, Toronto, ON, Canada, ⁴Dentist

ABSTRACT

Background: Atherosclerosis is the most common cause for heart attack and stroke. In the last decade, several epidemiological studies have found an association between periodontal infection and atherosclerosis. The aim of this research was to determine the possible association between chronic periodontal disease and severity of atherosclerosis.

Materials and Methods: Eighty-two subjects that were referred to Chamran Heart Hospital in Isfahan for angiography were involved in this study. Fifty-nine subjects had coronary artery obstruction (CAO) and 23 showed no obstruction after angiography. The severity of CAO was assessed. Periodontal parameters including pocket depth (PD), gingival recession (R), clinical attachment level (CAL), and bleeding on probing (BOP) of all subjects were recorded. The decayed-missing-filled (DMF) index of all subjects was also measured. For statistical analysis, Pearson correlation test, Chi-square, and independent *t*-test were used.

Results: There were significant positive correlation between variables R, PD, CAL, decayed (D), missing (M), DMF, BOP, and degree of CAO. However, there were no significant differences between filling variable degree of CAO (left anterior descending, left circumflex, and right coronary artery). Independent *t*-test showed that the mean of variables R, PD, AL, D, M, and DMF in patients with obstructed arteries were significantly higher than subjects without CAO. But there were no significant differences between variable F in two groups.

Conclusion: The results of this cross-section analytical study showed an association between periodontal disease and dental parameters with the severity of CAO measured by angiography. However, this association must not interpret as a cause and effect relationship.

Key Words: Angiography, association, coronary artery obstruction, periodontal disease

Received: April 2015
Accepted: December 2015

Address for correspondence:
Dr. Mohammad Ketabi,
Department of Periodontics,
Faculty of Dentistry,
Isalmic Azad University,
Isfahan (Khorasgan) Branch,
Isfahan, Iran.
E-mail: ketabimohammad@
yahoo.com

INTRODUCTION

In the last two decades, the strong association between periodontitis and many systemic conditions^[1] such as premature delivery,^[2] low weight birth infants,^[3] metabolic syndrome,^[4] and particularly cardiovascular diseases such as

myocardial infarction (MI)^[5] and coronary heart diseases (CHDs)^[6] have been shown.

In this regard, epidemiologic data and meta-analyses have suggested an association between periodontal disease and cardiovascular diseases.^[7-11] Among

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Ketabi M, Meybodi FR, Asgari MR. The association between periodontal disease parameters and severity of atherosclerosis. *Dent Res J* 2016;13:250-5.

Access this article online



Website: www.drj.ir
www.drjjournal.net
www.ncbi.nlm.nih.gov/pmc/journals/1480

cardiovascular diseases, atherosclerosis is the most common cause for heart attack and stroke. Most commonly, people develop atherosclerosis as a result of diabetes, genetic risk factors, high blood pressure, a high-fat diet, obesity, high blood cholesterol levels, and smoking. However, a sizable number of patients suffering from atherosclerosis do not harbor the classical risk factors. Ongoing infections like periodontal disease have been suggested to play a role in this process. Bartova *et al.*^[12] found an association between periodontal infection and atherosclerosis in particular. Obviously, the ultimate and dangerous result of atherosclerosis would be the partial or complete obstruction of coronary arteries. Hyvärinen *et al.*^[13] considered periodontal infection as a potential risk factor for cardiovascular disease and, in particular, atherosclerosis.

The number and degree of coronary artery obstruction (CAO) are usually measured by angiography. In angiography, the commonly number of obstructed branches namely left anterior descending (LAD), left circumflex (LCX), and right coronary artery (RCA) and their degree of obstructions are measured. Although, the association between periodontal disease and cardiovascular diseases are well established, but scarce information exists regarding the relation between periodontal disease with extend and severity of CAO. The aim of this research was to determine the possible association between periodontal (clinical attachment level [CAL], pocket depth [PD], bleeding on probing [BOP], R) and dental (D, M, F, decayed-missing-filled [DMF]) parameters with degree and number of obstructed coronary arteries measured by angiography.

MATERIALS AND METHODS

One hundred and fifty subjects that were referred to Chamran Heart Hospital in Isfahan for angiography were involved in this study. Patients with any systemic disease like diabetes, smokers, body mass index of more than 25 and patient with refractory periodontitis were excluded. Finally, 82 patients aged mean 55.08 (40–70) were recruited for the study. Out of 82 included subjects 58 had chronic periodontitis. After performing angiography, 59 had some degree of CAO but 23 had no CAO. The severity of CAO were assessed by 2 factors namely the number of obstructed main coronary arteries (1, 2 or 3,

i.e., LAD, LCX, RCA) and degree of obstruction. Periodontal parameters including PD, R, CAL, and BOP of all subjects were recorded. The DMF index of all subjects was also measured. For PD measurement, standard Williams's periodontal probe was applied using the blind method in 6 sites of every tooth. Patients were grouped as periodontitis if they had 3 pockets with the depth of more than 3 mm. O'Leary microbial plaque index was used for recording the superficial plaques (mesial, distal, lingual, and buccal). At the examination time, a dental disclosing tablet was given to the patients. Having patients chewed the tablet and washed of the mouth, dyed surfaces were examined using the probe tip to find the soft accumulations in the dentogingival junction. For BOP, the Ainamo and Bay index (1975) based on which the probe sulcus twisted was used. Any bleeding after 10 s was considered as the BOP and recorded in the examination chart. Gingival recession was recorded in mm and CAL was calculated as the sum of PD and R at each site.

Statistical analysis

To determine the association of numerical (quantitative) variables with each other, Pearson correlation test and for qualitative variables, Chi-square tests were used. For evaluation of the association of qualitative variables with quantitative variables independent *t*-test used.

RESULTS

Table 1 shows the general, periodontal, dental and cardiac characteristic of study population:

The mean age of subjects was 55.08 (44–69). Out of 82 subjects, 58 had chronic periodontitis (28% mild, 57% moderate, and 15% severe). After performing angiography, 59 subjects had some degree of CAO, but 29 had no CAO. The details of dental and periodontal variables and percentage of CAOs are also given.

Table 2 shows the frequency distribution and percentage obstruction of LAD, LCX, and RCA in all subjects.

Table 3 shows the correlation coefficient of R, PD, AL, D, M, F, DMF, and BOP with the percentage of LAD, LCX, and RCA obstruction. The Pearson correlation coefficient showed that there was a significant positive correlation between variables R, PD, AL, D, M, DMF, BOP, and the degree of coronary artery (LAD, LCX,

Table 1: Characteristic of study population including details of the dental and periodontal findings with individual age and cardiac parameters

Subjects with and without CAO and status of periodontal and dental conditions	Numbers and Means
Number of subject	82
CAO	59
No CAO	23
Status of periodontal condition	
Chronic periodontitis	58
Generalized gingivitis	21
Healthy gingiva	9
CAL (mm)	2.62±1.62
PPD (mm)	2.84±1 (number: 59 - obstructed artery) 0.52±0.67 (number: 23 - nonobstructed artery)
BOP	45.2%±12.5
R (mm)	1.2±0.64 (number: 59 - obstructed artery) 0.39±0.5 (number: 23 - nonobstructed artery)
D	4.93±3.23
M	8.6±4.75
F	3.89±3.09
DMF	17.43±7.23
LAD obstruction percentage	54.53±45.45
LCX obstruction percentage	43.79±44.85
RCA obstruction percentage	41.35±44.88
Obstructed arteries (number)	1.47±1.13
Number of patients with obstructed arteries	59
Number of patients without obstructed arteries	23

DMF: Decayed-missing-filled; RCA: Right coronary artery; LCX: Left circumflex; LAD: Left anterior descending; BOP: Bleeding on probing; PPD: Probing pocket depth; CAL: Clinical attachment level; CAO: Coronary artery obstruction; D: Decayed; M: Missing; F: Filled; R: Recession

Table 2: Frequency distribution and percentage of obstruction of left anterior descending, left circumflex and right coronary artery in subjects

Percentage of obstruction	LAD		LCX		RCA	
	n	Percentage	n	Percentage	n	Percentage
0-50	35	42.7	43	52.4	48	58.5
51-70	3	3.7	3	3.7	1	1.2
71-90	17	20.7	17	20.7	11	13.4
91-100	27	32.9	19	23.2	22	26.9
Total	82	100	82	100	82	100

LAD: Left anterior descending; LCX: Left circumflex; RCA: Right coronary artery

RCA) obstruction. However, there were no significant differences between variable F and CAO (LAD, LCX, and RCA).

Table 4 shows the mean of variables R, AL, D, M, F, and DMF in patients with and without obstructed arteries. Independent *t*-test showed that the mean of variables R, PD, AL, D, M, and DMF in patients

Table 3: Correlation coefficient of recession, pocket depth, clinical attachment level, decayed, missing, filled, decayed-missing-filled, bleeding on probing with percentage of left anterior descending, left circumflex and right coronary artery obstruction

Variables	LDA		LCX		RCA	
	r	P	r	P	r	P
R	0.523	<0.001	0.489	<0.001	0.433	<0.001
PD	0.533	<0.001	0.589	<0.001	0.627	<0.001
CAL	0.607	<0.001	0.632	<0.001	0.635	<0.001
D	0.500	<0.001	0.515	<0.001	0.495	<0.001
M	0.468	<0.001	0.674	<0.001	0.669	<0.001
F	0.027	0.405	-0.063	0.286	-0.026	0.286
DMF	0.543	<0.001	0.646	<0.001	0.650	<0.001
BOP	0.574	<0.001	0.638	<0.001	0.575	<0.001

D: Decayed; M: Missing; F: Filled; R: Recession; BOP: Bleeding on probing; PD: Pocket depth; CAL: Clinical attachment level; LCX: Left circumflex; RCA: Right coronary artery; DMF: Decayed-missing-filled

Table 4: The mean of variables recession, clinical attachment level, decayed, missing, filled, decayed-missing-filled in patients with and without obstructed arteries

Variables	Nonobstructed arteries (number of subjects: 23)		Obstructed arteries (number of subjects: 59)		P
	Mean	SD	Mean	SD	
R	0.39	0.5	1.2	0.64	<0.001
PPD	0.52	0.67	2.08	1	<0.001
CAL	0.91	0.9	3.29	1.33	<0.001
D	1.83	1.47	6.15	2.91	<0.001
M	3.87	2.05	10.46	4.19	<0.001
F	3.17	2.76	4.17	3.19	0.192
DMF	8.87	3.4	20.78	5.32	<0.001

D: Decayed; M: Missing; F: Filled; R: Recession; DMF: Decayed-missing-filled; CAL: Clinical attachment level; PPD: Probing pocket depth; SD: Standard deviation

with obstructed arteries were significantly higher than subjects without CAO. However, there was no significant difference between variable F in two groups.

DISCUSSION

The present study was a cross-sectional investigation of periodontal and dental parameters including PD, R, BOP, DMF, M, D, and F in patients with CAO (undergone for angiography). An association between these parameters (except filling variable) and number and degree of CAO were assessed. The results showed a strong association between parameters of periodontal disease with number and degree of CAO. These results are similar to other studies which

have proven periodontal infection as a potential risk factor for cardiovascular disease and, in particular, atherosclerosis.^[6,13]

However, a cross-sectional study from Sweden by Frisk *et al.*^[14] reported no such relation between dental infections and cardiovascular disease; however, their study population was all female and looked at only endodontic variables and CHD and did not look at periodontal variables.

The American Heart Association supports the hypothesis that there is an association between periodontitis and atherosclerotic vascular disease, independent of their confounders.^[15]

Few studies have also shown the association between extent and severity of periodontitis with extent and severity of cardiovascular disease (CVD). Kodovazenis *et al.*^[16] in a case-control study showed a consistent association between periodontitis and acute MI (AMI). The strength of the association increased concomitantly with the robustness of the criteria used to define periodontitis.

Holmlund and Lind^[17] could show in their investigation that the number of missing teeth was inversely correlated with the number of carotid arteries with atherosclerotic plaques.

Marfil-Álvarez *et al.*^[5] in a cross-sectional and analytical study reported the first research data demonstrating that the extent and severity of periodontitis is positively associated with AMI size (in patients who underwent diagnostic coronary angiography) as measured by serum troponin I and myoglobin levels.

Also to the best of our knowledge, this is the only study looked at association of periodontal infection and extend and severity of CAO measured by coronary angiography.

There are following possible explanations for this association:

- The role of periodontal pathogens: Hyvärinen *et al.*^[13] reported significantly higher salivary levels of *Aggregatibacter actinomycetemcomitans* in patients with stable coronary artery disease (CAD) compare to subjects showing no pathological findings by coronary angiography. High salivary levels of *A. actinomycetemcomitans* and systemic exposure to the bacterium were associated with increased risk for CAD. In this regard, biological evidence suggests a potential

causal role for periodontal bacteria that could be involved directly in atherogenesis^[18,19] or indirectly by increasing circulating cytokines and inflammatory mediators. Periodontal bacteria could migrate from the gingival/dental sites to the vascular wall via the bloodstream and act directly via their virulence factors, such as gingipains, proteinases, fimbriae, and/or lipopolysaccharides. In addition, the capacity of periodontal bacteria to induce leukocyte recruitment may contribute to this biological process.^[20] Numerous clinical studies have shown the presence of DNA from periodontal pathogens in atherosclerotic plaques^[21] and others have recovered viable, *Porphyromonas gingivalis* and *A. actinomycetemcomitans* from plaque samples^[22-24]

- In CHD patients with periodontitis, BOP is strongly associated with some strong systemic inflammatory mediators like C-reactive protein (CRP) levels; this association possibly reflects the potential significance of the local periodontal inflammatory burden for systemic inflammation^[25]
- Endothelium damage caused by the formation of lipid stripes may lead to bacteria penetrating into blood circulation after oral cavity procedures for patients with aggressive and chronic periodontitis^[13]
- The possibility of a potential role for periodontal microorganisms, especially *T. forsythia*, in neutrophil activation within hemorrhagic atherosclerotic carotid plaques has also been shown.^[20] The results of this and many other studies indicate a strong association between periodontal infection and CHD. On the other hand, interventional studies demonstrated beneficial effects of scaling and root planning on systemic levels of CRP, fibrinogen, and white blood cells.^[26] Therefore, a proper preventive and therapeutic program for periodontal diseases would be essential for these patients. Reichert *et al.*^[27] showed that use of floss/interdental brushes was associated with lower risk for new cardiovascular events among patients with CHD. Saffi *et al.*^[28] also showed that periodontal therapy lowered the level of CRP and other proinflammatory biomarkers in patients with stable CAD.

CONCLUSION

There was a clear association between periodontal disease parameter like PD, R, BOP, and obstruction of coronary arteries measured by Angiography. Therefore,

a proper treatment and good periodontal maintenance program could have a positive effect on the stability, severity, and extent of CHD. A well-controlled study which possibly compares the progression of CHD in periodontitis patients with and without a therapeutic periodontal program is suggested.

However, this was an analytical study with cross-section design, and its results could only establish an association between periodontal and oral parameters with CAO but this must not interpret as a causative or merely cause and effect relationship.

ACKNOWLEDGMENT

Authors would like to thank Mrs. Yaghmaei for her dedicated support and help to this manuscript.

Financial support and sponsorship
Nil.

Conflicts of interest

The authors of this manuscript declare that they have no conflicts of interest, real or perceived, financial or non-financial in this article.

REFERENCES

- Chrysanthakopoulos NA, Chrysanthakopoulos PA. Association between indices of clinically-defined periodontitis and self-reported history of systemic disorders in Greece. *Oral Health Prev Dent* 2013;11(3):251-60.
- Mathew RJ, Bose A, Prasad JH, Muliylil JP, Singh D. Maternal periodontal disease as a significant risk factor for low birth weight in pregnant women attending a secondary care hospital in South India: A case-control study. *Indian J Dent Res* 2014;25:742-7.
- Khadem N, Rahmani ME, Sanaei A, Afiat M. Association between preterm and low-birth weight with periodontal disease: A case-control study. *Iran J Reprod Med* 2012;10:561-6.
- Gurav AN. The association of periodontitis and metabolic syndrome. *Dent Res J (Isfahan)* 2014;11:1-10.
- Marfil-Álvarez R, Mesa F, Arrebola-Moreno A, Ramírez-Hernández JA, Magán-Fernández A, O'Valle F, *et al.* Acute myocardial infarct size is related to periodontitis extent and severity. *J Dent Res* 2014;93:993-8.
- Heaton B, Applebaum KM, Rothman KJ, Brooks DR, Heeren T, Dietrich T, *et al.* The influence of prevalent cohort bias in the association between periodontal disease progression and incident coronary heart disease. *Ann Epidemiol* 2014;24:741-6.
- Blaizot A, Vergnes JN, Nuwwareh S, Amar J, Sixou M. Periodontal diseases and cardiovascular events: Meta-analysis of observational studies. *Int Dent J* 2009;59:197-209.
- Humphrey LL, Fu R, Buckley DI, Freeman M, Helfand M. Periodontal disease and coronary heart disease incidence: A systematic review and meta-analysis. *J Gen Intern Med* 2008;23:2079-86.
- Bahekar AA, Singh S, Saha S, Molnar J, Arora R. The prevalence and incidence of coronary heart disease is significantly increased in periodontitis: A meta-analysis. *Am Heart J* 2007;154:830-7.
- Mustapha IZ, Debrey S, Oladubu M, Ugarte R. Markers of systemic bacterial exposure in periodontal disease and cardiovascular disease risk: A systematic review and meta-analysis. *J Periodontol* 2007;78:2289-302.
- Khader YS, Albashaireh ZS, Alomari MA. Periodontal diseases and the risk of coronary heart and cerebrovascular diseases: A meta-analysis. *J Periodontol* 2004;75:1046-53.
- Bartova J, Sommerova P, Lyuya-Mi Y, Mysak J, Prochazkova J, Duskova J, *et al.* Periodontitis as a risk factor of atherosclerosis. *J Immunol Res* 2014;2014:636893.
- Hyvärinen K, Mäntylä P, Buhlin K, Paju S, Nieminen MS, Sinisalo J, *et al.* A common periodontal pathogen has an adverse association with both acute and stable coronary artery disease. *Atherosclerosis* 2012;223:478-84.
- Frisk F, Hakeberg M, Ahlqwist M, Bengtsson C. Endodontic variables and coronary heart disease. *Acta Odontol Scand* 2003;61:257-62.
- Lockhart PB, Bolger AF, Papananou PN, Osinbowale O, Trevisan M, Levison ME, *et al.* Periodontal disease and atherosclerotic vascular disease: Does the evidence support an independent association? A scientific statement from the American Heart Association. *Circulation* 2012;125:2520-44.
- Kodovazenitis G, Pitsavos C, Papadimitriou L, Vrotsos IA, Stefanadis C, Madianos PN. Association between periodontitis and acute myocardial infarction: A case-control study of a nondiabetic population. *J Periodontal Res* 2014;49:246-52.
- Holmlund A, Lind L. Number of teeth is related to atherosclerotic plaque in the carotid arteries in an elderly population. *J Periodontol* 2012;83:287-91.
- Chukkapalli SS, Rivera MF, Velsko IM, Lee JY, Chen H, Zheng D, *et al.* Invasion of oral and aortic tissues by oral spirochete *Treponema denticola* in ApoE(-/-) mice causally links periodontal disease and atherosclerosis. *Infect Immun* 2014;82:1959-67.
- Velsko IM, Chukkapalli SS, Rivera MF, Lee JY, Chen H, Zheng D, *et al.* Active invasion of oral and aortic tissues by *Porphyromonas gingivalis* in mice causally links periodontitis and atherosclerosis. *PLoS One* 2014;9:e97811.
- Rangé H, Labreuche J, Louedec L, Rondeau P, Planesse C, Sebbag U, *et al.* Periodontal bacteria in human carotid atherothrombosis as a potential trigger for neutrophil activation. *Atherosclerosis* 2014;236:448-55.
- Reyes L, Herrera D, Kozarov E, Roldán S, Progulske-Fox A. Periodontal bacterial invasion and infection: Contribution to atherosclerotic pathology. *J Clin Periodontol* 2013;40 Suppl 14:S30-50.
- Kozarov EV, Dorn BR, Shelburne CE, Dunn WA Jr., Progulske-Fox A. Human atherosclerotic plaque contains viable invasive *Actinobacillus actinomycescomitans* and *Porphyromonas gingivalis*. *Arterioscler Thromb Vasc Biol* 2005;25:e17-8.
- Li L, Michel R, Cohen J, Decarlo A, Kozarov E. Intracellular survival and vascular cell-to-cell transmission of *Porphyromonas gingivalis*. *BMC Microbiol* 2008;8:26.
- Rafferty B, Jönsson D, Kalachikov S, Demmer RT, Nowygrod R, Elkind MS, *et al.* Impact of monocytic cells on recovery of

- uncultivable bacteria from atherosclerotic lesions. *J Intern Med* 2011;270:273-80.
25. Bokhari SA, Khan AA, Butt AK, Hanif M, Izhar M, Tatakis DN, *et al.* Periodontitis in coronary heart disease patients: Strong association between bleeding on probing and systemic biomarkers. *J Clin Periodontol* 2014;41:1048-54.
 26. Bokhari SA, Khan AA, Butt AK, Azhar M, Hanif M, Izhar M, *et al.* Non-surgical periodontal therapy reduces coronary heart disease risk markers: A randomized controlled trial. *J Clin Periodontol* 2012;39:1065-74.
 27. Reichert S, Schlitt A, Beschow V, Lutze A, Lischewski S, Seifert T, *et al.* Use of floss/interdental brushes is associated with lower risk for new cardiovascular events among patients with coronary heart disease. *J Periodontol Res* 2015;50:180-8.
 28. Saffi MA, Furtado MV, Montenegro MM, Ribeiro IW, Kampits C, Rabelo-Silva ER, *et al.* The effect of periodontal therapy on C-reactive protein, endothelial function, lipids and proinflammatory biomarkers in patients with stable coronary artery disease: Study protocol for a randomized controlled trial. *Trials* 2013;14:283.

